

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/002755

International filing date: 15 March 2005 (15.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/553,494
Filing date: 16 March 2004 (16.03.2004)

Date of receipt at the International Bureau: 22 April 2005 (22.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

PA 1285808

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office


February 23, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/553,494

FILING DATE: March 16, 2004

**By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS**



**M. SIAS
Certifying Officer**

13146 U.S. PTO

Docket Number 70436

FILING BY "EXPRESS MAIL" UNDER 37 CFR §1.10

EV 035650134 US
Express Mail Label Number

March 16, 2004
Date of Deposit

22264 U.S. PTO
60/553494



Address to: Commissioner for Patents
Box Provisional Patent Application
P.O. Box 1450
Alexandria, VA 22313-1450

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

Transmitted herewith for filing under 37 CFR §1.53(c) is the PROVISIONAL APPLICATION for patent of

INVENTOR(S)		
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
Stefan	BAUM	Muenchwilten, Switzerland
TITLE OF THE INVENTION (280 characters max) PESTICIDAL COMPOSITIONS		

CORRESPONDENCE ADDRESS
Direct all correspondence to the address associated with Customer No. 0026748, which is currently: Syngenta Crop Protection, Inc. Patent and Trademark Dept. 410 Swing Road Greensboro, NC 27409

ENCLOSED APPLICATION PARTS (check all that apply)
<input checked="" type="checkbox"/> Specification (Including Any Claims and Abstract) - ²⁴ pages <input type="checkbox"/> Drawings - sheets <input type="checkbox"/> Other (specify):

METHOD OF PAYMENT	
The Commissioner is hereby authorized to charge filing fee and any additional fees required to Deposit Account Number: 50-1676 in the name of Syngenta Crop Protection, Inc.	PROVISIONAL FILING FEE AMOUNT: \$ 160

☐ U.S. Government agency and contract number: (if the invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.)

Respectfully submitted,

William A. Teoli, Jr.

William A. Teoli, Jr.
Attorney for Applicants
Reg. No. 33,104
Tel. No. (336) 632-7706

Date: March 16, 2004

The PTO did not receive the following listed item(s) page 22 of 24
specification

Pesticidal Compositions

The present invention relates to a pesticidal composition and a slurry composition comprising the pesticidal composition, which compositions are particularly useful in treating propagation material to protect it against attack by pests; to a pest resistant propagation material; to a method of protecting propagation material; to a method of controlling pests from crop plants; and to use of certain surface active compounds for improving characteristics of the treated material.

Treatment of propagation material with pesticidal compositions allows protection against soil-borne pests during a susceptible stage in its development. Further, systemic seed treatments may provide an alternative to the traditional foliar pesticidal applications. However, following such treatment, the treated propagation material is handled and conveyed for packing and storage; this handling and conveying often results in the treated material brushing against each other to cause the chemicals that were bound to the material to become unbound. The release of the chemicals is often in the form of air-borne particles of, for example, pesticides and other materials from the treating composition that on drying the seed form solid particles. This can result in the treated material losing its pesticidal effectiveness and exposing workers in such treatment operations to health risks, through inhalation and sensitisation, for example, eye irritancy, of the air-borne particles.

The challenge is to improve the adherence of such particles to the propagation material.

It has been found that a pesticidal composition comprising certain type of surface active compounds in a defined ratio provides a reduction in the air-borne particles (commonly referred to as 'dust') when the treated propagation material, such as a seed, is handled (improved dust-off).

Accordingly, in a first aspect, the present invention provides a pesticidal composition, preferably a seed treating composition, such as a seed coating composition, comprising:

- (A) at least one pesticide, and optionally at least one other substance which has a melting point above 30°C; and
- (B) at least two surface active compounds, wherein (i) at least one surface active compound has a molecular weight of less than 2200, preferably less than 1700, such as in the range 400 to 1500, advantageously in the range 600 to 1200, and a Hydrophile-Lipophilic Balance (HLB) of at least 10, preferably in the range 10 to 25, such as 12 to 20, preferably 14 to 18

and (ii) at least one surface active compound is non-ionic, has a molecular weight of at least 2200, preferably at least 3000, such as in the range of 3500 to 15000, for example, 3500 to 10000, especially 4000 to 7500, advantageously 4500 to 6000 where the hydrophile constituent of the compound contributes to 10 to 85, such as 15 to 80, preferably 17 to 50, % of the molecular weight and, independent of the hydrophile proportion, the molecular weight of the hydrophobe constituent of the compound is from 2000 to 10000, preferably 2400 to 3900, more preferably 3000 to 3800, such as 3200 to 3700;

provided that the weight ratio of (B):(A) is in the range 0.08 to 0.5, preferably 0.1 to 0.3, advantageously 0.15 to 0.25, and the weight ratio of (ii):(i) is at least 0.5, such as at least 1.0, preferably at least 1.5, especially in the range 2 to 5, advantageously in the range 2 to 3.

In a second aspect, the present invention provides a slurry composition comprising the pesticidal composition of the first aspect, a liquid carrier and optionally (i) one or more, preferably other, formulation adjuvants, (ii) one or more other pesticidal compositions, each comprising at least one further pesticide, or both (i) and (ii).

In a third aspect, the present invention provides a method of protecting propagation material, preferably a seed, from attack by pests, such as soil-dwelling pests, preferably a nematode, by treating the material with a pesticidal, preferably a nematocidal, effective amount of the composition of the first or second aspect, preferably before planting or sowing the seed.

Accordingly, the invention also provides a method controlling pests, such as a nematode, from damaging crop plants, especially vegetable crop plants, by the treatment of the plant's propagation material, such as a seed, with a with a pesticidal, preferably a nematocidal, effective amount of the composition of the first or second aspect, preferably before planting or sowing the seed.

In a fourth aspect, the present invention provides a pest resistant plant propagation material comprising a plant propagation material, preferably a seed, such as a vegetable seed, treated with a pesticidal, preferably nematocidal, effective amount of the composition of the first or second aspect or obtained by the method of the third aspect.

In a fifth aspect, the present invention provides a use in a pesticidal composition or slurry composition, to improve the dust-off property of a propagation material, preferably a seed, that has been treated with the composition, of:

- 5 at least two surface active compounds, wherein (i) at least one surface active compound has a molecular weight of less than 2200 and a Hydrophile-Lipophilic Balance (HLB) of at least 10 and (ii) at least one surface active compound is non-ionic, has a molecular weight of at least 2200 where the hydrophile constituent of the compound contributes to 10 to 85 % of the molecular weight and the molecular weight of the hydrophobe constituent of the
10 compound is from 2000 to 10000;

provided that the weight ratio of the surface active compounds to one or more pesticides, and optionally at least one other substance which has a melting point above 30°C, in the pesticidal composition or slurry composition, is in the range 0.08 to 0.5, and the weight ratio
15 of (ii):(i) is at least 0.5.

- In another aspect, the present invention provides a method of improving the dust-off property of a propagation material, preferably a seed, that has been treated with a pesticidal or slurry composition, said method comprising mixing in the pesticidal or slurry composition
20 at least two surface active compounds, wherein (i) at least one surface active compound has a molecular weight of less than 2200 and a Hydrophile-Lipophilic Balance (HLB) of at least 10 and (ii) at least one surface active compound is non-ionic, has a molecular weight of at least 2200 where the hydrophile constituent of the compound contributes to 10 to 85 % of the molecular weight and the molecular weight of the hydrophobe constituent of the
25 compound is from 2000 to 10000;

provided that the weight ratio of the surface active compounds to one or more pesticides, and optionally at least one other substance which has a melting point above 30°C, in the pesticidal composition or slurry composition, is in the range 0.08 to 0.5, and the weight ratio
30 of (ii):(i) is at least 0.5.

The invention is described in more detail below.

Compositions

- 35 After seed treatment, the substances that can become air-borne particles have a melting point above 30°C, such substances are pesticides, colouring agents, thickening agents.

In one embodiment, the pesticide (or active ingredient) is present in the composition of the first aspect in an amount of from about 12.5 % to about 50 % by weight, more specifically, from 30 to about 45% by weight of the composition; the balance of the composition, also known as a formulation, comprising a solid and/or liquid carrier along with surfactant(s) and other optional inert ingredients known in the art as formulation adjuvants, e.g., protective colloids, adhesives, thickeners, thixotropic agents, penetrating agents, preservatives, stabilizers, antifoaming agents, antifreeze agents, sequestering agents, colourings, such as dyes or pigments, and polymers.

In the composition of the second aspect, the proportions of the components would be less than that in the first aspect and depends on the amount of the liquid carrier, which is normally present in a major proportion, and also on the presence of one or more, of the same or, other formulation adjuvants and/ or one or more other pesticidal compositions each comprising a further pesticide.

The pesticide can be of any type, for example, a fungicide, such as triazole derivatives, strobilurins, carbamate (including thiocarbamate), benzimidazoles (thiabendazole), N-trihalomethylthio compounds (captan), substituted benzenes, carboxamides, phenylamides and phenylpyrroles, and mixtures thereof; an insecticide (such as neonicotinoids, carbamates and pyrethroids), acaricide, molluscicide and a nematocide.

Preferred examples of suitable pesticides for a composition of the first aspect or second aspect are selected from abamectin (1), acephate (2), acetamiprid (4), alpha-cypermethrin (202), azinphos-methyl (45), bifenthrin (76), carbaryl (115), carbofuran (118), carbosulfan (119), chlorpyrifos (145), clothianidin (165), cyromazine (209), deltamethrin (223), dimethoate (262), emamectin benzoate (291), endosulfan (294), fipronil (354), furathiocarb (412), gamma-HCH (430), imidacloprid (458), Isufenphos, methiocarb (530), omethoate (594), tefluthrin (769), thiamethoxam (792), thiacloprid (791), thiodicarb (799), azoxystrobin (47), pyraclostrobin (690), benomyl (62), bitertanol (84), captan (114), carbendazim (116), carboxin (120), chlorothalonil (142), copper salts (such as copper sulfate (172), cuprous oxide (181), Bordeaux mixture (87), copper hydroxide (169), copper sulfate (tribasic) (173), copper oxychloride (171) and copper octanoate (170)), cymoxanil (200), cyproconazole (207), cyprodinil (208), difenoconazole (247), diniconazole (267), ethirimol, famoxadone (322), fenamidone (325), fenhexamid (334), fenpiclonil (341), fluazinam (363), fludioxonil (368), fluquinconazole (385), flutolanil (396), flutriafol (397), fosetyl-aluminium (407), fuberidazole (409), guazatine (422), hexaconazole (435), hymexazol (447), imazalil (449), iprodione (470), isufenphos, mancozeb (496), maneb (497), metalaxyl (516), metalaxyl-M

(517), metconazole (525), myclobutanil (564), silthiofam (729), nuarimol (587), oxadixyl (601), oxine-copper (605), oxolinic acid (606), pencycuron (620), prochloraz (659), procymidone (660), pyrimethanil (705), pyroquilon (710), quintozone (716), tebuconazole (761), tetraconazole (778), thiabendazole (790), thiophanate-methyl (802), thiram (804), triadimenol (815), triazoxide (821), triticonazole (842), trifloxystrobin (832), picoxystrobin (647) and ipconazole (468).

In an embodiment of either the first or second aspect, the pesticide is selected from the active ingredients abamectin (1), emamectin benzoate (291), chlorpyrifos (145), metalxyl-M (517), thiamethoxam (792), difenoconazole (247), azoxystrobin (47), tefluthrin (769), fludioxnil (368), imidacloprid (458), thiacloprid (791), fipronil (354), clothianidin (165), thiram (804) and myclobutanil (564).

Especialmente, a composição do primeiro aspecto compreende Abamectin (1), como o pesticida, e, vantajosamente, a composição do segundo aspecto compreende, como pesticidas, abamectin (1), emamectin benzoato (291), clorpirifos (145), tiametoxam (792), tefluthrin (769), fludioxnil (368), imidacloprid (458), tiacloprid (791), fipronil (354), clothianidin (165) e thiram (804).

The pesticides are described in the e-Pesticide Manual, version 3.0, 13th Edition, Ed. CDC Tomlin, British Crop Protection Council, 2003-04. The number following the compound name is the entry number given in the Pesticide Manual.

Compositions of the first aspect come in a variety of formulations: dry flowables (DF), liquid flowables (LF), true liquids (TL), emulsifiable concentrates (EC), suspension concentrates (SC), dusts (D), wettable powders (WP), suspoemulsions (SE), water-dispersible granules (WG) and others. Some are registered for use only by commercial applicators using closed application systems, others are readily available for on-farm use as dusts, slurries, water-soluble bags, or liquid ready-to-apply formulations.

Preferably, the composition according to the first aspect or second aspect is an aqueous composition, for example, a two-phase system, such as an emulsion and a suspension aqueous concentrates, preferably in the form of suspension concentrate. Suspension concentrate formulations, for example, for seed treatment, generally have a viscosity of 300 to 1200, such as 400 to 800, mPas⁻¹, when measured in a BROOKFIELD viscometer with spindle 3 at 30 rpm and 25°C. The average size of the suspended particles is 0.1 to 20,

especially 0.5 to 5, advantageously 1 to 3, microns, when measured with a laser particle analyzer, e.g., a Malvern Mastersizer S.

5 The formulations, *i.e.*, the compositions, preparations or combinations containing (A) and (B) components of the first aspect, as well as, if appropriate, suitable inert solid or liquid carriers, are prepared in a known manner, for example, by intimately mixing and/or grinding the components with inert, agriculturally-acceptable extenders, for example with solid or liquid carriers.

10 The term "carrier" in the present description denotes a natural or synthetic, organic or inorganic material with which the active ingredient, *i.e.*, the pesticide, is combined in order to facilitate its application to the seeds. This carrier is hence generally inert, and it must be agriculturally acceptable, in particular to the seed being treated. The carrier may be solid (clays, natural or synthetic silicates, silica, resins, waxes, solid fertilizers, and the like) or
15 liquid (water, alcohols, ketones, petroleum fractions, aromatic or paraffinic hydrocarbons, chlorinated hydrocarbons, liquefied gases, and the like).

Suitable liquid carriers are: aromatic hydrocarbons, in particular the fractions C₈ to C₁₂, such as xylene mixtures or substituted naphthalenes, phthalic esters such as dibutyl or dioctyl
20 phthalate, aliphatic hydrocarbons such as cyclohexane or paraffins, alcohols and glycols as well as their ethers and esters, such as ethylene glycol monomethyl ether, ketones such as cyclohexanone, strongly polar solvents such as N-methyl-2-pyrrolidone, dimethyl sulfoxide or dimethylformamide, and, if appropriate, epoxidized vegetable oils or soybean oil; or
25 water.

A preferred carrier for compositions according to the present invention (both the first and second aspects) is a liquid carrier, advantageously water.

30 The amount of surface active compounds (B) generally present range from 1 to 25, preferably 2.4 to 22.5, especially 5 to 10, %, by weight, based on the weight of the composition of the first aspect. Surface active compounds are made up of water soluble (hydrophilic) components (or constituents), such as polyoxyethylene, and water insoluble (hydrophobic) components (or constituents), such as polyoxypropylene. Examples of
35 surface active compounds are surfactants having good emulsifying, dispersing and wetting properties, depending on the nature of the pesticide to be formulated. Surfactants will also be understood as meaning mixtures of surfactants. Surfactants are non-ionic, cationic and/or anionic.

The (B)(i) surface active compound preferably has a molecular weight of at least 100.

The (B)(ii) surface active compound preferably has a molecular weight of at most of 100,000.

In an embodiment, the weight ratio of surface active compounds (ii) to (i) is at most 10.

In an embodiment, two surface active compounds (B)(ii) are present in the composition.

In the event, two surface active compounds (B)(ii) are present:

- the first surface active compound has a molecular weight of the hydrophobe constituent of from 2400 to 3900, preferably 3000 to 3800, such as 3200 to 3700 and, independent of the hydrophobe molecular weight, a proportion of the molecular weight of the hydrophile constituent of from 13 to 45, preferably 17 to 40, such as 18 to 30, %; and
- the second surface active compound has a molecular weight of the hydrophobe constituent of from 2200 to 3900, preferably 2500 to 3600, such as 2700 to 3200 and, independent of the hydrophobe molecular weight, a proportion of the molecular weight of the hydrophile constituent of from 43 to 67, preferably 45 to 65, such as 50 to 60, %.

The Hydrophile-Lipophilic Balance (HLB) value is an index of the hydrophilic nature of a compound proposed by Griffin. The HLB value of a polyoxyethylene alkyl ether can be determined by, for example, the Griffin equation.

HLB value = [(molecular weight of the hydrophilic moiety)/(molecular weight of the surface active compound)] x 20

Groups, for example, sulfate and phosphate ions, can also contribute to the HLB value.

Generally, compounds, including surface active compounds, that are commercially used tend to be not analytically pure, but a mixture of suitable compounds, for example, of the same chemistry but of different analogs, isomers and molecular weights. The characteristics attributed to, for example, the (B)(i) and (B)(ii) surface active compounds are, therefore, preferably also satisfied in a mixture of compounds where the characteristics are possessed by a compound in the mixture, which compound is present in a major proportion, such as greater than 50, preferably greater than 60, especially greater than 75,

% by weight, based on the weight of the mixture; more preferably, the mixture itself satisfies the characteristics defined.

5 The surfactants customarily employed in formulation technology are described, *inter alia*, in the following publications:

"McCutcheon's Detergents and Emulsifiers Annual", MC Publishing Corp., Glen Rock, N.J., 1988.

10 M. and J. Ash, "Encyclopedia of Surfactants", Vol. I-III, Chemical Publishing Co., New York, 1980-1981.

15 Among the suitable surfactants there may be mentioned, *e.g.*, polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or (mono- or di-alkyl)naphthalenesulphonic acid salts, polystyrene sulfonate salts, laurylsulfate salts, polycondensates of ethylene oxide with
20 lignosulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols such as mono- and di-(polyoxyalkylene alkylphenol) phosphates, polyoxyalkylene alkylphenol carboxylates or polyoxyalkylene alkylphenol sulfates), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyltaurides), polycondensates of ethylene
oxide with phosphated tristyrylphenols and polycondensates of ethylene oxide with phosphoric esters of alcohols or phenols.

25 In an embodiment, the (B)(i) surface active compound is an ionic, advantageously an anionic, surfactant; preferably one or more (B)(i) surface active compound is selected from a sulfate type (*e.g.*, an aryl sulfate) and a phosphate type (such as an alkylphenol polyalkoxyether phosphate, a block copolymer of polyalkoxyether phosphate, polyarylphenol polyalkoxyether phosphate and an arylphenol polyalkoxyether phosphate),
30 especially a phosphate type surfactant (such as a polyarylphenol polyalkoxyether phosphate). Particularly desired in the compositions of the invention are that each (B)(i) surface active compound is of the same type, a preferred type is a phosphate type surfactant.

35 The anionic surfactants may be present as acids or include alkali metals (such as lithium, sodium and potassium), alkali earth metals (such as calcium and magnesium), ammonium and various amines (such as alkylamines, cycloalkylamines and alkanolamines).

Specific examples of suitable anionic surfactants include: Sorprophor PS19 (Rhodia), Dowafax 30 C05 (Dow), Soprophor 4D384 (Rhodia) and Soprophor 3D33 (Rhodia).

5 In an embodiment, the (B)(ii) surface active compound is a polyalkylene oxide polymer, such as a block polymer. Specific examples are polyoxyethylene polyoxypropylene block polymers, and polyoxyethylene polyoxypropylene block polymer ethers. Specific examples include Toximul 8320 (Stepan), Emulsogen 3510 (Clariant), AntaroX PL/122 (Rhodia), Pluronic L101 (BASF), Pluronic L122 (BASF) and Pluronic PE 10500 (BASF).

10 The compositions of the invention can also contain a wetting agent, which is also considered surface active compound in that it has a water soluble (hydrophilic) and water insoluble (hydrophobic) components, but they tend to non-ionic and generally have a molecular weight of less than 2000, and so can be a component according to (B)(i). In a preferred embodiment of the first aspect, a wedding agent is not present.

15 Advantageously, three surface active compounds, one of (B)(i) and two of (B)(ii), are used in the compositions according to the first aspect.

20 The compositions of the invention also may comprise at least one antifreeze agent. In an embodiment, the antifreeze agent is present in at least about 2 and up to about 25% of, more specifically from 3 to about 10% by weight, based on the weight of the composition of the first aspect.

25 Specific examples of suitable antifreezes include ethylene glycol, 1,2-propylene glycol, 1,3-propylene glycol, 1,2-butanediol, 1,3-butanediol, 1,4-butanediol, 1,4-pentanediol, 3-methyl-1,5-pentanediol, 2,3-dimethyl-2,3-butanediol, trimethylol propane, mannitol, sorbitol, glycerol, pentaerythritol, 1,4-cyclohexanedimethanol, xlenol, bisphenols such as bisphenol A or the like. In addition, ether alcohols such as diethylene glycol, triethylene glycol, tetraethylene glycol, polyoxyethylene or polyoxypropylene glycols of molecular weight up to
30 about 4000, diethylene glycol monomethylether, diethylene glycol monoethylether, triethylene glycol monomethylether, butoxyethanol, butylene glycol monobutylether, dipentaerythritol, tripentaerythritol, tetrapentaerythritol, diglycerol, triglycerol, tetraglycerol, pentaglycerol, hexaglycerol, heptaglycerol, and octaglycerol.

35 As a particularly preferred subset of suitable antifreeze materials there can be mentioned ethylene glycol, propylene glycol and glycerol.

The compositions of the invention optionally includes at least one polymer selected from water-soluble and water-dispersible film-forming polymers. Suitable polymers have an average molecular weight of at least about 7,000 up to about 200,000; more specifically at least about 10,000, up to about 100,000. The composition of the first aspect generally contain from about 0% to about 10% by weight of the composition of polymer. As used herein, "nonionic surfactants" are different compounds from the water-dispersible and water-soluble polymers described herein.

Suitable polymers are selected from

- d1) ethylene vinylacetate copolymers,
- d2) vinylacetate/vinylpyrrolidone copolymers,
- d3) alkylated vinylpyrrolidone copolymers,
- d4) polyvinylpyrrolidone, and
- d5) polyalkyleneglycol including the polypropylene glycols and polyethylene glycols.

The compositions of the invention also optionally contains at least one thickener. In one embodiment, the thickener is present in an amount from about 0.01% to about 5% w/w, more specifically from 0.05 to 2% by weight, based on the weight of the composition of the first aspect.

Illustrative of thickeners (water-soluble polymers which exhibit pseudoplastic properties in an aqueous medium) are gum arabic, gum karaya, gum tragacanth, guar gum, locust bean gum, xanthan gum, carrageenan, alginate salt, casein, dextran, pectin, agar, 2-hydroxyethyl starch, 2-aminoethyl starch, 2-hydroxyethyl cellulose, methyl cellulose, carboxymethyl cellulose salt, cellulose sulfate salt, polyacrylamide, alkali metal salts of the maleic anhydride copolymers, and alkali metal salts of poly(meth)acrylate.

As suitable thickeners there may also be mentioned attapulgate-type clay, carrageenan, croscarmellose sodium, furcelleran, glycerol, hydroxypropyl methylcellulose, polystyrene, hydroxypropyl cellulose, hydroxypropyl guar gum, and sodium carboxymethylcellulose. Xanthan gum and attapulgate-type clay are preferred.

The compositions of the invention can be employed together with the adjuvants customary in formulation technology, biocides, biostats, emulsifiers (Iethicin, sorbitan, and the like), antifoam agents or application-promoting adjuvants customarily employed in the art of formulation. In addition, there may be mentioned inoculants and brighteners.

Additionally, a colouring agent, such as a dye or pigment, is included in the seed coating so that an observer can immediately determine that the seeds are treated. The colouring agent is also useful to indicate to the user the degree of uniformity of the coating applied.

- 5 The compositions of the invention can be prepared by processes known in the art, such as forming a homogeneous suspension with all the components, except the thickeners, and wet milling the suspension until the desired particle size is reached, then the thickeners and further water are added to a set viscosity.
- 10 The final composition can be screened if desired to remove any insoluble particles.

In one embodiment, commercial products according to the first aspect will preferably be formulated as concentrates. They may be used undiluted or may be diluted with a liquid carrier, such as water, and one or more components. The decision to dilute (by adding
15 other components or liquid carrier) depends on the treatment methods available to the user. The composition according to the second aspect is an example of diluting the composition of the first aspect. The liquid carrier in the slurry composition tends to be the liquid carrier, if any, of the concentrate formulation of the first aspect.

- 20 The pesticidal composition of the first aspect may be used alone or in combination with other pesticidal compositions for treatment of propagation material either together or sequentially.

Generally, a composition containing a variety of components tend to interact with each other
25 and affect the physical characteristics of that composition, such as an increase in the sedimentation and a change in the viscosity over time and/ or climatic conditions of storage. Accordingly, compositions that exhibit good storage characteristics so that they can be transported long distances and kept in store-houses are also desired. It has been found that the composition of the first aspect also demonstrate such improved physical
30 characteristics.

Further, in an embodiment of the composition of the first aspect a polymer component is excluded from the composition because it has been found to affect the physical characteristics of the composition on storage. However, a polymer may be used in
35 combination with the composition of the first aspect in a slurry composition of the second aspect (e.g., tank mix) to provide additional coating properties before application to the propagation material because in such an event the resulting composition does not tend to

be stored and also, optionally, compatibility agents may be used. In the event a polymer is used in a slurry composition, the polymer is generally present in an amount of from 1 to 7, such as 2 to 6, % by weight, based on the weight of the composition of the first aspect.

- 5 Generally, a composition of the second aspect is that applied for controlling pests, such as by treating propagation material. In such an event, such a composition would contain more types of formulation components, than a composition of the first aspect, which is an ingredient in the preparation of the composition of the second aspect.

10 Uses

- The term "plant propagation material" is understood to denote all the generative parts of the plant, such as seeds, which can be used for the multiplication of the latter and vegetative plant material such as cuttings and tubers (for example, potatoes). There may be
- 15 mentioned, e.g., the seeds (in the strict sense), roots, fruits, tubers, bulbs, rhizomes, parts of plants. Germinated plants and young plants, which are to be transplanted after germination or after emergence from the soil, may also be mentioned. These young plants may be protected before transplantation by a total or partial treatment by immersion.
- 20 The pesticidal compositions according to first and second aspects are particularly suited to treatment of seeds. The techniques of seed treatment application are well known to those skilled in the art, and they may be used readily in the context of the present invention. For the purposes of this invention, seed treatments are defined as chemical or biological substances that are applied to seeds or vegetative plant propagation materials to control
- 25 disease organisms, insects, or other pests. The seed treatment composition includes the pesticides, such as fungicides, bactericides, nematocides and other classes of insecticides. Most seed treatments are applied to true seeds, which have a seed coat surrounding an embryo. However, some seed treatments can be applied to vegetative plant propagation materials such as rhizomes, bulbs, corms or tubers.
- 30 The composition of the first aspect may be used, for example, for treatment, in an undiluted form or be diluted with a liquid carrier, for example. In the instance it is diluted, such compositions are known as slurries or tank mix and represent an example of the second aspect of the invention. The compositions of the first and second aspect may be applied to a
- 35 seed to result in a film (or dressing), a coating, or a pellet, of the pesticide on the seed after drying, depending on the treatment process. The processes are well known in the art, and employ, for seeds, e.g., the techniques of film-coating or encapsulation, or for the other

propagation material, the techniques of, e.g., immersion. Needless to say, the method of application of the compounds to the seed may be varied and the invention is intended to include any technique which is to be used.

- 5 The compositions of the invention are formulated for protecting cultivated plants and their propagation materials. The compositions are advantageously formulated for seed treatment applications against soil inhabiting insects, which can damage the crop in the early stages of plant development. For example, the compositions can be formulated to target insects and representatives of the order Acarnia including:
- 10 from the order Lepidoptera, for example, *Acleris* spp., *Adoxophyes* spp., *Aegeria* spp., *Agrotis* spp., *Alabama argillaceae*, *Amylois* spp., *Anticarsia gemmatalis*, *Archips* spp., *Argyrotaenia* spp., *Autographa* spp., *Busseola fusca*, *Cadra cautella*, *Carposina nipponensis*, *Chilo* spp., *Choristoneura* spp., *Clysia ambiguella*, *Cnaphalocrocis* spp., *Cnephasia* spp., *Cochylis* spp., *Coleophora* spp., *Crocidolomia binotalis*, *Cryptophlebia leucotreta*, *Cydia* spp., *Diatraea* spp., *Diparopsis castanea*, *Earias* spp., *Ephestia* spp.,
- 15 *Eucosma* spp., *Eupoecilia ambiguella*, *Euproctis* spp., *Euxoa* spp., *Grapholita* spp., *Hedya nubiferana*, *Heliothis* spp., *Hellula undalis*, *Hyphantria cunea*, *Keiferia lycopersicella*, *Leucoptera scitella*, *Lithocollethis* spp., *Lobesia botrana*, *Lymantria* spp., *Lyonetia* spp., *Malacosoma* spp., *Mamestra brassicae*, *Manduca sexta*, *Operophtera* spp., *Ostrinia nubilalis*, *Pammene* spp., *Pandemis* spp., *Panolis flammea*, *Pectinophora gossypiella*, *Phthorimaea operculella*, *Pieris rapae*, *Pieris* spp., *Plutella xylostella*, *Prays* spp., *Scirpophaga* spp., *Sesamia* spp., *Sparganothis* spp., *Spodoptera* spp., *Synanthedon* spp., *Thaumetopoea* spp., *Tortrix* spp., *Trichoplusia ni* and *Yponomeuta* spp.;
- 20 from the order Coleoptera, for example,
- 25 *Agriotes* spp., *Anthonomus* spp., *Atomaria linearis*, *Chaetocnema tibialis*, *Cosmopolites* spp., *Curculio* spp., *Dermestes* spp., *Diabrotica* spp., *Epilachna* spp., *Eremnus* spp., *Leptinotarsa decemlineata*, *Lissorhoptrus* spp., *Melolontha* spp., *Orycaephilus* spp., *Otiorhynchus* spp., *Phlyctinus* spp., *Popillia* spp., *Psylliodes* spp., *Rhizopertha* spp., *Scarabeidae*, *Sitophilus* spp., *Sitotroga* spp., *Tenebrio* spp., *Tribolium* spp. and *Trogoderma* spp.;
- 30 from the order Orthoptera, for example,
- Blatta* spp., *Blattella* spp., *Gryllotalpa* spp., *Leucophaea maderae*, *Locusta* spp., *Periplaneta* spp. and *Schistocerca* spp.;
- from the order Isoptera, for example,
- 35 *Reticulitermes* spp.;
- from the order Psocoptera, for example,
- Liposcelis* spp.;

from the order Anoplura, for example,

Haematopinus spp., Linognathus spp., Pediculus spp., Pemphigus spp. and Phylloxera spp.;

from the order Mallophaga, for example,

5 Damalinae spp. and Trichodectes spp.;

from the order Thysanoptera, for example,

Frankliniella spp., Hercinothrips spp., Taeniothrips spp., Thrips palmi, Thrips tabaci and Scirtothrips aurantii;

from the order Heteroptera, for example,

10 Cimex spp., Distantiella theobroma, Dysdercus spp., Euchistus spp., Eurygaster spp., Leptocorisa spp., Nezara spp., Piesma spp., Rhodnius spp., Sahlbergella singularis, Scotinophara spp. and Triatoma spp.;

from the order Homoptera, for example,

Aleurothrixus floccosus, Aleyrodes brassicae, Aonidiella spp., Aphididae, Aphis spp.,
15 Aspidiotus spp., Bemisia tabaci, Ceroplaster spp., Chrysomphalus aonidium, Chrysomphalus dictyospermi, Coccus hesperidum, Empoasca spp., Eriosoma larigerum, Erythroneura spp., Gascardia spp., Laodelphax spp., Lecanium corni, Lepidosaphes spp., Macrosiphus spp., Myzus spp., Nephrotettix spp., Nilaparvata spp., Paratoria spp., Pemphigus spp., Planococcus spp., Pseudaulacaspis spp., Pseudococcus spp., Psylla
20 spp., Pulvinaria aethiopica, Quadraspidiotus spp., Rhopalosiphum spp., Saissetia spp., Scaphoideus spp., Schizaphis spp., Sitobion spp., Trialeurodes vaporariorum, Trioza erytreae and Unaspis citri;

from the order Hymenoptera, for example,

Acromyrmex, Atta spp., Cephus spp., Diprion spp., Diprionidae, Gilpinia polytoma,
25 Hoplocampa spp., Lasius spp., Monomorium pharaonis, Neodiprion spp., Solenopsis spp. and Vespa spp.;

from the order Diptera, for example,

Aedes spp., Antherigona soccata, Bibio hortulanus, Calliphora erythrocephala, Ceratitis spp., Chrysomyia spp., Culex spp., Cuterebra spp., Dacus spp., Drosophila melanogaster,
30 Fannia spp., Gastrophilus spp., Glossina spp., Hypoderma spp., Hyppobosca spp., Liriomyza spp., Lucilia spp., Melanagromyza spp., Musca spp., Oestrus spp., Orseolia spp., Oscinella frit, Pegomyia hyoscyami, Phorbia spp., Rhagoletis pomonella, Sciara spp., Stomoxys spp., Tabanus spp., Tannia spp. and Tipula spp.;

from the order Siphonaptera, for example, Ceratophyllus spp. und Xenopsylla cheopis and

35 from the order Thysanura, for example, Lepisma saccharina;
crucifer flea beetles (Phyllotreta spp.), root maggots (Delia spp.), cabbage seedpod weevil (Ceutorhynchus spp.) and aphids; and

from the order Acarina, for example, *Acarus siro*, *Aceria sheldoni*, *Aculus schlechtendali*, *Amblyomma* spp., *Argas* spp., *Boophilus* spp., *Brevipalpus* spp., *Bryobia praetiosa*, *Calipitimerus* spp., *Chorioptes* spp., *Dermanyssus gallinae*, *Eotetranychus carpini*, *Eriophyes* spp., *Hyalomma* spp., *Ixodes* spp., *Olygonychus pratensis*, *Ornithodoros* spp.,
5 *Panonychus* spp., *Phyllocoptruta oleivora*, *Polyphagotarsonemus latus*, *Psoroptes* spp.,
Rhipicephalus spp., *Rhizoglyphus* spp., *Sarcoptes* spp., *Tarsonemus* spp. and *Tetranychus* spp..

10 Protection of developing plants against attack by plant parasitic nematodes is also obtainable in the instance the pesticide is, for example, Abamectin. Abamectin is effective against a wide range of nematode pests including species of *Meloidogyne* (for example, *Meloidogyne incognita* and *Meloidogyne javanica*), *Heterodera* (for example, *Heterodera glycines*, *Heterodera schachtii*, *Heterodera avenae* and *Heterodera trifolii*), *Globodera* (for example, *Globodera rostochiensis*), *Radopholus* (for example, *Radopholus similis*),
15 *Rotylenchulus*, *Pratylenchus* (for example, *Pratylenchus neglectans* and *Pratylenchus penetrans*), *Aphelenchoides*, *Helicotylenchus*, *Hoplolaimus*, *Paratrichodorus* and *Tylenchorhynchus*, in particular *Meloidogyne*.

20 The composition of the first and second aspect, in the case abamectin is present as a pesticide, are particularly effective in controlling nematodes.

Optionally, in addition to the control of insect pests, the compositions of the invention advantageously are formulated with fungicides for seed treatment applications against diseases in the soil, which mostly occur in the early stages of plant development. For
25 example, the compositions can be formulated to target pathogens including *Pythium*, *Tilletia*, *Gerlachia*, *Septoria*, *Ustilago*, *Fusarium*, *Rhizoctonia* (so-called "damping off complex"); Oomycetes such as *Phytophthora*, *Plasmopara*, *Pseudoperonospora*, *Bremia* etc. as well as against the *Botrytis* species, *Pyrenophora*, *Monilinia* and further representatives of the *Ascomycetes*, *Deuteromycetes* and *Basidiomycetes* classes.
30

Suitable target crops are especially potatoes, cereals, (wheat, barley, rye, oats), rice, maize, sugar beet, cotton, millet varieties, sorghum, tobacco, sun flowers, beans, peas, oil plants (rape, canola), soybeans, cabbages, tomatoes, eggplants (aubergines), pepper and other vegetables, and spices as well as ornamental shrubs and flowers.
35

Suitable target crops also include transgenic crop plants of the foregoing varieties. The transgenic crop plants used according to the invention are plants, or propagation material

thereof, which are transformed by means of recombinant DNA technology in such a way that they are - for instance - capable of synthesizing selectively acting toxins as are known, for example, from toxin-producing invertebrates, especially of the phylum Arthropoda, as can be obtained from *Bacillus thuringiensis* strains; or as are known from plants, such as lectins; or in the alternative capable of expressing a herbicidal or fungicidal resistance. Examples of such toxins, or transgenic plants which are capable of synthesizing such toxins, have been disclosed, for example, in EP-A-0 374 753, WO 93/07278, WO 95/34656, EP-A-0 427 529 and EP-A-451 878 and are incorporated by reference in the present application.

The compositions are suited for dressing applications on plant propagation material. The latter term embraces seeds of all kinds (fruit, tubers, grains), cuttings, cut shoots and the like. The preferred field of application is the treatment of all kinds of seeds (as specified in the target crops above), and in particular the seed treatment of canola, maize, cereals, cotton, tomatoes, tobacco, soybeans, other legumes, and other vegetables and crops that are susceptible, especially preferred are vegetable crop seeds.

As noted above, the compositions of this invention may be formulated or mixed in the seed treater tank or combined on the seed by overcoating with other pelleting materials and/ or seed treating agents. The agents to be mixed with the compounds of this invention may be for the control of pests, nutrition, and the control of plant diseases.

The composition has particular application to concurrent (such as by slurry) and sequential seed treatments.

Seeds treated with the compositions of the invention generally have a drying time ranging from 20 to 60 seconds when being treated at room temperature. Generally, for vegetable seeds a spray drying technique is used.

The composition according to the first aspect contains and/or may be applied together or sequentially with further compounds on to the propagation material, such as a seed to form a seed coating or a pellet of the seed. These further compounds can be fertilizers or micronutrient donors or other preparations that influence plant growth. They can also be selective herbicides, fungicides, other insecticides, bactericides, insect growth regulators, plant growth regulators, nematocides, molluscicides or mixtures of several of these preparations.

Accordingly, a seed coating composition, particularly effective in the pelleting of seeds, is a preferred embodiment of the compositions of the first and second aspects.

5 A preferred method of applying the composition according to the invention consists in spraying or wetting the plant propagation material with the liquid preparation, or mixing the plant material with such liquid preparation. Also, before the application, the composition of the invention may be diluted with water by simple mixing at ambient temperature in order to prepare an on-farm seed treatment formulation.

10 The formulation may be applied, for low value crops, such as cereals, at application volumes ranging from 200ml to 3 liters per 100 kg seed, more specifically, from 400ml to 2 liters per 100 kg seed. For vegetable crop seeds, the amount tend to be higher.

15 A beneficial feature of the composition is that it provides an increased adherence of particles, in particular air-borne particles, to the seed, which results in decreased dustiness and the subsequent elimination of related dust problems. Elimination of the dust associated with many seed treatments also eliminates the associated health hazards to those who work with treated seeds, such as processing plant employees, truck drivers, warehouse workers, and farmers. The compositions of the present invention also enable satisfactory
20 flowability & individuality of the propagation material (*i.e.*, allow the material to be easily handled and not stick to each other).

Still another advantage of this invention is the uniform coating of seeds with non-dusting seed treatment which will not interfere with germination and sprouting of the seed but which
25 will protect the seed and resultant seedling against pests, particularly soil-borne pests.

The application of a the compositions according to the invention directly on a seed, seed piece or bare root dip treatment for early season crop protection has several distinct advantages both from an economic and environmental standpoint. By treating seeds, seed
30 pieces or bare roots, preferably prior to planting or sowing, with a pesticide, for example, a nematocide in the case of Abamectin, the pesticide is concentrated at the site of pest, *e.g.*, nematode, attack and, therefore, much less active ingredient on a per acre basis is required in comparison with conventional soil application. This not only makes pest control more economic, but also minimizes environmental disturbance and any adverse non-target
35 effects.

As used in the present application, the embodiments of the features of the present invention are independent of each other, unless otherwise specified.

The following Examples are given by way of illustration and not by way of limitation of the invention.

EXAMPLES

Examples A, B, C and 1 to 13 are prepared by mixing surfactants, neutraliser (if any), a defoaming agent, an antifreeze agent, and a bactericide with water until a homogeneous phase is achieved. Subsequently, a colour pigment (if any) and active ingredient are added and are mixed. The resulting mixture is then wet-milled through a so-called bead mill (Dyno, Drais, Premier for instance). The milling parameters are set in such a way that the average particle size of the resulting ground premix is within specifications (usually median particle size average below 1.5um). Finally, the polymer (if any), thickening agent and a minor amount of water are added and the final product is mixed for at least 30 minutes. Table 1 indicates their compositional details, with each composition containing identical amounts of defoaming agent, thickeners and bactericide. The numbers given in the Table are weight.

The formulations are treated onto wheat seeds by Hege seed treatment device and then the amount of dust given off is measured by placing the treated seeds in a steel drum with blades, which when the drum is rotated stimulates handling and conveying of the seeds. A precision airflow control system provides a constant flow of air that carries air borne particles through a coarse filter separator onto a filter. The dust quantity is measured by weighing the filter (results are tabulated in Table 1).

Table 1:

Examples																
Component	A	B	1	2	3	4	C	5	6	7	8	9	10	11	12	13
Abamectin (85%)	47.00	47.00	47.00	47.00	47.00	47.00	40.80	47.00	47.00	47.00	47.00	47.00	47.00	42.50	42.50	40.80
anionic surfactant, dispersant		2.00					3.00									3.00
anionic surfactant, dispersant	2.00		2.00	2.00	2.00	2.00		2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	
conv. wetting agent, surfactant	1.00	1.00	1.00	1.00	1.00	1.00	2.00	1.00	1.00					1.00	1.00	2.00
polymeric surfactant, dispersant																
polymeric surfactant, dispersant																
polymeric surfactant, dispersant			2.00	2.00												
polymeric surfactant, dispersant																
polymeric surfactant, dispersant	1.00	1.00			2.00	2.00			4.00		6.00		6.00	6.00		
neutralizer	0.20	0.20	0.20	0.20	0.20	0.20		0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	5.00
antifreeze	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	5.00	5.00	5.00	5.00	5.00	5.00	4.00
colour pigment										0.50			0.50	0.50	0.50	
60%polymerdispersion in water			4.00				5.00									
60%polymerdispersion in water				4.00	4.00											
water	52.80	52.80	47.80	47.80	47.80	51.80	53.20	49.80	49.80	47.30	47.80	47.80	47.30	50.80	50.80	53.20
(A) - Abamectin & other solids*	40.9	40.9	40.9	40.9	40.9	40.9	35.4	40.9	40.9	41.4	40.9	40.9	41.4	37.5	37.5	35.4
(B) - total surfactant	4.0	4.0	5.0	5.0	5.0	5.0	5.0	7.0	7.0	8.0	8.0	8.0	8.0	9.0	9.0	10.0
surfactant (i)	3.00	3.00	3.00	3.00	3.00	3.00	5.00	3.00	3.00	2.00	2.00	2.00	2.00	3.00	3.00	5.00
surfactant (ii)	1.00	1.00	2.00	2.00	2.00	2.00	0.00	4.00	4.00	6.00	6.00	6.00	6.00	6.00	6.00	5.00
ratio of B / A	0.10	0.10	0.12	0.12	0.12	0.12	0.14	0.17	0.17	0.19	0.20	0.20	0.19	0.24	0.24	0.28
ratio of (ii) / (i)	0.33	0.33	0.67	0.67	0.67	0.67	0.00	1.33	1.33	3.00	3.00	3.00	3.00	2.00	2.00	1.00
Dust-off measurements	17.8 g	22.4 g	12.80 g	12.40 g	12.20 g	12.90 g	20.5 g	12.9 g	10.60 g	8.90 g	8.7 g	9.9 g	10.20 g	5.20 g	5.80 g	2.8 g

* these materials are considered, on drying of the seeds, to form solid particles on the seed

February 23, 2005

410 SWING ROAD
GREENSBORO, NC 27409

This is to inform you that the document requested in your order dated 02-10-2005 is a true reproduction of the official office record copy of that document:

60553494 CERTIFIED PAT APP AS FILED-MAIL 10

PAGE 20 IS MISSING FROM THE SPECIFICATION.

☒ The enclosed office document is a reproduction from the official office record copy and has been recorded using high quality scanning or microfilm equipment. Copies of page/papers that were not scannable have not been included, nor have pages/papers received after the original filing date. Copies of these pages/papers may be ordered separately.

☐ The enclosed document is a reproduction of the best available source of the official office record copy of that document.

If you have any questions or need additional information, please contact our Customer Service Department.

Mailing Address:

Mail Stop Document Services,
Director of the U.S. Patent and Trademark Office
P.O. Box 1450
Alexandria, VA 22313-1450

Delivery Address:

U.S. Patent and Trademark Office
Office of Public Records, Customer Service
1213 Jefferson Davis Highway, Suite 300
Arlington VA 22202

Order your copies of USPTO documents on the Web. It's fast, easy, and secure! Go to (<http://ebiz1.uspto.gov/oems25p/index.html>) and order today!

Voice: (703) 308-9726

Fax: (703) 308-7048

E-Mail: <mailto:DSD@USPTO.GOV>

Ref:MS 1285808

CLAIMS

1. A pesticidal composition comprising:

(A) at least one pesticide, and optionally at least one other substance which has a melting point above 30°C; and

(B) at least two surface active compounds, wherein (i) at least one surface active compound has a molecular weight of less than 2200 and a Hydrophile-Lipophilic Balance (HLB) of at least 10 and (ii) at least one surface active compound is non-ionic, has a molecular weight of at least 2200 where the hydrophile constituent of the compound contributes to 10 to 85 % of the molecular weight and the molecular weight of the hydrophobe constituent of the compound is from 2000 to 10000;

provided that the weight ratio of (B):(A) is in the range 0.08 to 0.5, and the weight ratio of (ii):(i) is at least 0.5.

2. The composition of claim 1, wherein the ratio of (B):(A) is 0.1 to 0.3, preferably 0.15 to 0.25.

3. The composition of either claim 1 or claim 2, wherein the ratio of surface active compounds (ii):(i) is at least 1.0, preferable at least 1.5, especially in the range 2 to 5, advantageously in the range 2 to 3.

4. The composition according to any one of claims 1 to 3, wherein (B)(i) is ionic, preferably anionic.

5. The composition according to any one of claims 1 to 4, wherein (B)(i) has a molecular weight of less than 1700, such as in the range 400 to 1500, preferably in the range 600 to 1200.

6. The composition according to any one of claims 1 to 5, wherein (B)(i) has a HLB value in the range 10 to 25, such as 12 to 20, preferably 14 to 18.

7. The composition according to any one of claims 1 to 6, wherein (B)(ii) has a molecular weight of at least 3000, such as in the range of 3500 to 15000, especially 4000 to 7500, advantageously 4500 to 6000.

8. The composition according to any one of claims 1 to 6, wherein two surface active compounds (B)(ii) are present.
- 5 9. The composition according to claim 7, wherein the first surface active compound has a molecular weight of the hydrophobe constituent of from 2400 to 3900, preferably 3000 to 3800, such as 3200 to 3700 and, independent of the hydrophobe molecular weight, a proportion of the molecular weight of the hydrophile constituent of from 13 to 45, preferably 17 to 40, such as 18 to 30, %; and the second surface active compound has a molecular
10 weight of the hydrophobe constituent of from 2200 to 3900, preferably 2500 to 3600, such as 2700 to 3200 and, independent of the hydrophobe molecular weight, a proportion of the molecular weight of the hydrophile constituent of from 43 to 67, preferably 45 to 65, such as 50 to 60, %.
- 15 10. The composition according to any one of claims 1 to 9, wherein a (B)(i) surface active compound is selected from a sulfate type surfactant and a phosphate type surfactant, especially a phosphate type surfactant.
- 20 11. The composition according to any one of claims 1 to 10, wherein each (B)(ii) surface active compound is a polyalkylene oxide polymer.
12. The composition according to claim 11, where each copolymer is a block polymer.
- 25 13. The composition according to any one of claims 1 to 12, wherein (A) the pesticide is selected from abamectin, emamectin benzoate, chlorpyrifos, metalxyl-M, thiamethoxam, difenoconazole, azoxystrobin, tefluthrin, fludioxnil, imidacloprid, thiacloprid, fipronil, clothianidin, thiram and myclobutanil.
- 30 14. A slurry composition comprising the pesticidal composition according to any one of claims 1 to 13, a liquid carrier and optionally (i) one or more formulation adjuvants, (ii) one or more other pesticidal compositions, each comprising at least one further pesticide, or both (i) and (ii).
- 35 15. A method of protecting propagation material from attack by pests by treating the material with a pesticidal effective amount of the composition claimed in any one of claims 1 to 14.

16 A pest resistant plant propagation material comprising a plant propagation material treated with a pesticidal effective amount of the composition claimed in any one of claims 1 to 14, or obtained by the method claimed in claim 15.

5 17. Use in a pesticidal composition or slurry composition, to improve the dust-off property of a propagation material that has been treated with the composition, of:

10 at least two surface active compounds, wherein (i) at least one surface active compound has a molecular weight of less than 2200 and a Hydrophile-Lipophilic Balance (HLB) of at least 10 and (ii) at least one surface active compound is non-ionic, has a molecular weight of at least 2200 where the hydrophile constituent of the compound contributes to 10 to 85 % of the molecular weight and the molecular weight of the hydrophobe constituent of the compound is from 2000 to 10000;

15 provided that the weight ratio of the surface active compounds to one or more pesticides, and optionally at least one other substance which has a melting point above 30°C, in the pesticidal composition or slurry composition, is in the range 0.08 to 0.5, and the weight ratio of (ii):(i) is at least 0.5.

20

ABSTRACT

Pesticidal Compositions

- 5 A pesticidal composition comprising:
- (A) at least one pesticide, and optionally at least one other substance which has a melting point above 30°C; and
- 10 (B) at least two surface active compounds, wherein (i) at least one surface active compound has a molecular weight of less than 2200 and a Hydrophile-Lipophilic Balance (HLB) of at least 10 and (ii) at least one surface active compound is non-ionic, has a molecular weight of at least 2200 where the hydrophile constituent of the compound contributes to 10 to 85 % of the molecular weight and the molecular weight of the hydrophobe constituent of the compound is from 2000 to 10000;
- 15 provided that the weight ratio of (B):(A) is in the range 0.08 to 0.5, and the weight ratio of (ii):(i) is at least 0.5. Such compositions demonstrate improved dust-off performance when applied to propagation material, such as seeds.